

Preventative Care Checklist

Children (birth to 10 years)



Discuss your checklist with your primary care provider at your annual wellness visit.
CHI Saint Joseph Medical Group — Primary Care | 859.313.2255

Well Child Exams and Vaccines		
COVID-19	6 months and older	
Exam and Hepatitis B	Birth to 10 years	
Exam	1 month	
Exam, Pentacel, Prevnar, Hepatitis B and Rotavirus	2 months	
Exam, Pentacel, Prevnar and Rotavirus	4 months	
Exam, Pentacel, Prevnar, Hepatitis B, Flu Shot and Rotavirus	6 months	
Exam	9 months	
Exam, Anemia Test, Possible TB and Lead Tests, MMR, Varicella and Hepatitis A	12 months	
Exam, Pentacel, and Prevnar	15 months	
Exam, Flu Shot and Hepatitis A	18 months	
Exam	24 months	
Exam and Flu Shot	3 years	
Exam, Flu Shot, DTaP, Polio, MMR, and Varicella	4 years	
Exam and Flu Shot	5 years	
Exam and Flu Shot	6 - 10 years (every 1 - 2 years)	
Depression and Suicide Risk	12 years and older	
Screenings		
Body Mass Index	2 years and older	
Dental Caries	Birth - 5 years	
Hepatitis B	Birth - 10 years	
Obesity	6 years and older	
Skin Cancer Behavioral Counseling	6 months - 10 years (with fair skin)	
Vision	3 - 5 years	

*The information in these checklists include recommendations adapted from the following sources as of January 2023 and is subject to change. Preventative services with a rating of A or B from the U.S. Preventive Services Task Force. Immunizations for children, adolescents and adults recommended by the Centers for Disease Control; and preventative care screenings for infants, children, adolescents and women supported by the Health Resources and Services Administration. Based on risk factors and as advised by your health care provider.

Immunization Requirements for Preschool - 12th Grade

3 Years	4 Years	5 Years	11-15 Years	16 Years and Older
DTaP - 4 doses	*DTaP - 4 or 5 doses	*DTaP - 4 or 5 doses	*DTaP - 4 or 5 doses	*DTaP - 4 or 5 doses
Polio (IPV) - 3 doses	*Polio (IPV) - 3 or 4 doses	*Polio (IPV) - 3 or 4 doses	*Polio (IPV) - 3 or 4 doses	*Polio (IPV) - 3 or 4 doses
MMR - 1 dose	MMR - 2 doses	MMR - 2 doses	MMR - 2 doses	MMR - 2 doses
*Hib - 3 or 4 doses	*Hib - 3 or 4 doses	*Hib - no requirement	Tdap - 1 dose	Tdap - 1 dose
*HepA - 2 doses effective for the school year beginning on or after July 1, 2018	*HepA - 2 doses effective for the school year beginning on or after July 1, 2018	*HepA - 2 doses effective for the school year beginning on or after July 1, 2018	*HepA - 2 doses effective for the school year beginning on or after July 1, 2018	*HepA - 2 doses effective for the school year beginning on or after July 1, 2018
HepB - 3 doses	HepB - 3 doses	HepB - 3 doses	HepB - 3 doses	HepB - 3 doses
*Varicella - 1 dose	*Varicella - 2 doses	*Varicella - 2 doses	*Varicella - 2 doses	*Varicella - 2 doses
*Pneumococcal (PCV) - 4 doses	*Pneumococcal (PCV) - 4 doses	*Pneumococcal (PCV) - no requirement	Meningococcal ACWY (MCV4) - 1 dose (at 11 years of age)	Meningococcal ACWY (MCV4) - 2 doses for 16 yrs and older effective for the school year beginning on or after July 1, 2018

*DTaP - Fifth dose not needed if dose #4 given after 4th B-day and 6 months from previous dose

*Polio - Fourth dose not needed if dose #3 given after 4th B-Day and 6 months from previous dose

*Hib - last dose must be on or after 12 months of age - not required at age 5 and older; number of total doses (3 or 4) depends on brand used

*PCV - last dose must be on or after 12 months of age - not required at age 5 years and older

*Varicella - no doses needed if history of Chickenpox or Zoster disease is verified by a medical provider

*Meningococcal ACYW - if initial dose is given at 16 yrs or older, will only need one dose (previously known as MCV4)

*Hepatitis A - 2 doses spaced 6 months apart effective for the school year beginning on or after **July 1, 2018**